

SOME TOPICS OF GEODYNAMIC EVOLUTION OF INNER WESTERN CARPATHIANS

Z. NÉMETH¹ and M. PUTIŠ²

¹*Geological Survey of Slovak Republic, Jesenského 8, 040 01 Košice, Slovakia; nemeth@gssr-ke.sk*

²*Dpt. of Mineralogy and Petrology, Faculty of Science, Comenius Univ., Mlynská dol., 842 15 Bratislava, Slovakia;
putis@fns.uniba.sk*

Abstract: Finding of a Variscan geosuture in the North-Gemeric zone modified former interpretations of Variscan tectonic evolution of Inner Western Carpathians. South-vergent ductile shear zone between Early Paleozoic Gelnica and Rakovec Groups of Gemericum manifested kinematics of Variscan exhumation. We suppose that thermal processes in collisional régime were sufficient for increase of thermal gradient, metamorphism and metallogenic processes.

Key words: geosuture, geodynamics, Gelnica and Rakovec Groups, Gemericum

The long-lasting research of Gemericum and surrounding units, including detail geological mapping, structural research in meso/microscales and accounting of published results of further branches of geo-sciences, brought some new topics into interpretations of geodynamic evolution of Inner Western Carpathians.

Variscan geodynamic evolution

Variscan evolution terminated with origin of *Rakovec geosuture* including lithology of Rakovec and Klátov Groups and partly Carboniferous rocks. From the Lower Paleozoic Gelnica Group it is separated with ductile shear zone of moderate northern inclination. This inclination of rock boundaries in suture zone was proved also by the results of the project SGR-Geophysics (Grecula and Kuchariè, eds., 1992) as well as by the series of seismic profiles G1a/92, G1b/92 and G2/93 (Vozár et al., 1998). Microtectonic research of mylonites from the boundary zone between Gelnica and Rakovec Groups (Németh, 2001; Németh et al., 2001a) proved two principal directions of tectonic transport - in autochthonous footwall of ductile shear zone (that is in upper part of Gelnica Group) the general vergency of tectonic

transport is ESE. In allochthonous hanging wall of shear zone (tectonic *mélange* of Rakovec Group) the transport directions are generally to the south, locally with wider azimuthal spread. Our findings support the overall southern vergency of Variscan tectogenesis (Putiš, 1992; Putiš and Grečula in Plašienka et al., 1997). South-vergent structures are common in the Gemicum, encompassing fold structures.

The Lower Paleozoic rock sequences of Gemicum from the viewpoint of evolution can be generally divided into two large groups:

A. The rocks of Gelnica Group, representing the Lower Paleozoic cover of rigid blocks of older crystalline basement of Panafrican? provenience, were in Upper Carboniferous deformed by exhumed *mélange* of rocks of Rakovec Group (+ Klátov Group + part of Carboniferous sequences).

B. Rocks of Rakovec Group suffered tectonometamorphism in various depths of subduction zone. We suppose the Devonian-Westphalian age of subduction-exhumation processes. The dating of the upper boundary is based only on finding of the pressure peak of 12 kbars in a part of Westphalian sediments of Rudòany Formation (Radvanec, 1998). It suggests that also Westphalian sediments were locally recrystallized in subduction régime and subsequently exhumed. Subduction slab was inclined northwards below rocks of recent Veporicum and Tatricum. Subduction initiated the reverse displacement of rock masses in subduction slab back to surface (including sporadical mantle fragments; cf. Radvanec, 1999). When the crystalline rocks of southern basement (underlying Gelnica Group) had approached the subduction zone, the convergence decelerated and stopped.

Collision resetted the thermal régime in collided terrane and the deeply entrenched isograds neighbouring the subduction canal started to be straightened. The thermal régime during Permian was highly probable related also to the presence of prolonged thermal source below the collided terrane in the North-Gemic zone. Position of thermal axis in North-Gemic zone results also from reconstruction of the whole Carpathian Permian sedimentary basin and Permian magmatic trends (Vozárová and Vozár, 1987). We suppose that the increased thermal flow caused also higher temperature metamorphism of rock sequences in Gelnica Group, origin of Gemic granites and metallogenic processes.

Later Permian evolution in the Inner Western Carpathians is characterized with northward displacement of lithospheric plates with collisional terrane. The position of the axis of conductive heat was therefore gradually shifted towards the southernmore Gemic zones. This is proven also by still younger ages of Gemic granite from the north southward:

[Hnilec - 290 Ma (Kovach et al., 1986) and 282 Ma (Cambel et al., 1989); Betliar 272 Ma; Humel 270 and 246 Ma; Zlatá Idka 251 and 223 Ma (Kovach et al., 1986)]. Similar effect is exhibited also by the presence of best developed siderite-sulphidic veins directly in North-Gemic zone. These veins penetrate from the Gelnica Group through the suture zone. Southward the veins parameters become worse. Conductive overheating of Lower Paleozoic rocks of the Gelnica Group was most intensive in its lowermost horizons, built with black schists, lydites, carbonates and the variegated volcanic horizon (Betliar Fm., Holec Beds and lower parts of Smolník Fm.). Rocks served as a source of metals for fluids entering into fractures in overlying rock sequences (cf. Grecula et al., 1991). Position of mentioned rocks on rigid crystalline basement and below soft cover of overlying rocks of Gelnica Group protected them against tectonization during preceding south-vergent exhumation of rocks of the Rakovec Group. Pre-Permian tectogenesis was accommodated by overlying rocks of Smolník and Hnilec Fms. (Fig. 1).

Alpine geodynamic evolution

The effects of Permian extension in Gemic terrane, owing to displacement of lithosphere above the axis of convectional heat, were shifted southwards. The continental crust in South-Gemic zone was gradually disintegrated and the prolonged Meliata-Hallstatt basin originated. Later Alpine evolution after Upper Jurassic? convergence has been divided into four deformation stages:

Deformation stage AD₁ is tightly related to the closure of Meliata-Hallstatt basin with following succession of events: Closure of the basin and transport of the Bôrka nappe on Gemicum. Ongoing convergence caused not only the north-vergent imbrication of Lower Paleozoic rocks of former Variscan collisional terrane, but the Lower Cretaceous displacement of several kilometers thick rock pile of Gemicum northward on Veporicum. This process of detachment was allowed by the rheologically contrast horizon of the black schists of Betliar Fm., dividing rigid crystalline basement from relatively softer rocks of Gelnica Group. We suppose that a part of former Betliar Formation remained in its homeland. The north-vergent imbrication in low-temperature brittle-ductile and brittle regime caused multiple repeating the Early Paleozoic lithology of Gemicum in recent picture. Thrusts destructed also courses of Late-Variscan ore veins (cf. numerous examples in Grecula et al., 1995).

The Lower Cretaceous Gemic nappe displacement during AD₁ caused:

1. Truncation of northern depth continuation of Rakovec geosuture, and "retraction" of its lower parts below Gemicum.

2. Thrusting of North-Gemic zone on "Veporic ramp" caused uplift of frontal parts of the nappe, its disintegration and later erosive removal of cover carbonates of the Stratená Group.

Thickening of continental crust caused the Middle Cretaceous unroofing of Veporic "core" during AD₂. The younger conjugate systems of brittle-ductile and brittle shear zones of NW-SE and NE-SW trends (AD₃; cf. Grecula et al., 1990; Németh et al., 2001b) formed arc bending of Gemicum, and *sensu lato* of the whole Western Carpathians.

The orogen parallel extension in east-west direction during AD₄, documented preferably in the North-Gemic zone, was caused by crustal indentor directed to the north to the Gemic zone.

References

- Cambel, B., Bagdasaryan, G. P., Gukasyan, R. C. & Veselský, J., 1989: Rb-Sr geochronology of leucocratic granitoid rocks from the Spišsko-gemerské rudohorie Mts. and Veporicum. *Geol. Zbor. Geol. carpath.*, 40, 323-332.
- Cambel, B., Kráľ, J. & Burchart, J., 1990: Isotopic geochronology of crystalline basement in Western Carpathians. *Veda, Bratislava*, 183. (In Slovak).
- Grecula, P., Návesňák, D., Bartalský, B., Gazdačko, L., Németh, Z., Ištván, J. & Vrbatovič, P., 1990: Shear zones and arc structure of Gemicum, the Western Carpathians. *Mineralia Slov.*, 22, 97-110.
- Grecula, P., Radvanec, M. & Bartalský, B., 1991: Critical thermic isograde in metamorphic-hydrothermal model of vein mineralization on the background of the Variscan events, Gemic unit, Western Carpathians. *Mineralia Slov.*, 23, 403-411.
- Grecula, P. & Kucharič, J., red., 1992: Partial final report from comprehensive geological-geophysical interpretation of the northern part of SGR. Manuscript, archives ŠGÚDŠ, Bratislava, 199. (In Slovak)
- Grecula, P. et al., 1995: Mineral deposits of Slovak Ore Mountains. Vol. 1. Geocomplex, Bratislava, 1-829.
- Kováč, A., Svingor, E. & Grecula, P., 1986: Rb-Sr isotopic ages of granitoid rocks from the Spišsko-gemerské rudohorie Mts., Western Carpathians, Eastern Slovakia. *Mineralia Slov.*, 18, 1-14.
- Németh, Z., 2001: Petrotectonics of the ductile shear zones of Gemicum. Ph.D. thesis. Comenius Univ., Bratislava, 1-98.
- Németh, Z., Grecula, P. & Putiš, M., 2001a: Lithotectonic relations in boundary zone of Gelnicka and Rakovec Group in the North-Gemic zone. *Geol. práce, Správy* 105, 67-70. (In Slovak)
- Németh, Z., Putiš, M. & Grecula, P., 2001b: Origin of the arc bended boundary of Gemicum and Veporicum from the viewpoint of kinematics of Alpine unroofing. *Geol. práce, Správy* 105, 65-66. (In Slovak)
- Plašienka, D., Grecula, P., Putiš, M., Kováč, M. & Hovorka, D., 1997: Evolution and structure of the Western Carpathians: an overview. In: Grecula, P., Hovorka, D. & Putiš, M. (Eds.): *Geological evolution of the Western Carpathians. Mineralia Slov. - Monograph*, Bratislava, 1-24i.
- Putiš, M., 1992: Variscan and Alpidic nappe structures of the Western Carpathian crystalline basement. *Geol. Carpath.* 43, 6, 369-380.
- Radvanec, M., 1998: High-pressure metamorphism of Upper Carboniferous conglomerate from the locality Rudňany-Svinský hrb on the north of Gemicum. *Mineralia Slov.*, 30, 95-108. (In Slovak)
- Radvanec, M., 1999: Eclogitized clinopyroxenic gabbro with retrograde metamorphism in pumpellyite-actinolite facies on the hills Babiná and Ostrá (Gemicum). *Mineralia Slov.*, 31, 467-484. (In Slovak)
- Vozár, J., Szalaoiová, V. & Šantavý, J., 1998: Interpretation of the Western Carpathian deep structures on the basis of gravimetric and seismic sections. In: M. Rakús, ed.: *Geodynamic development of the Western Carpathians. Dionýz Štúr Publishers*, Bratislava, 241-257.

Vozárová, A. & Vozár, J., 1987: West Carpathians Late Paleozoic and its paleotectonic development. In: H. W. Flügel, F. P. Sassi & P. Grech (eds.): Pre-Variscan and Variscan events in the Alpine-Mediterranean mountain belts. Mineralia Slov. - Monograph, Alfa, Bratislava, 469-487.

Fig. 1. Interpretation of the geodynamics of Inner Western Carpathians. GE - Gelnica Group, RA - Rakovec Group, KL - Klátov Group, C - Carboniferous rocks, P - Permian rocks. White arrows - direction of convergence, resp. divergence, black undulating arrows - axis of convective heat, black straight arrows - relative migration of collided terrane with respect of axis of convective heat. 1 - black schists of Betliar Fm. (Ordovician-Silurian), 2 - green schists of Smolník Fm. (S-D), 3 - flysch - lower bed - in Smolník Fm. (S-D), upper bed - the supreme parts of Hnilec Fm. (D-C₁?), 4 - intermediate and acid volcanoclastics of Hnilec Fm. (D), 5 - effusive and extrusive products of basalt volcanism, locally with accompanying rocks (D), 6 - clastics of Rudňany Fm. (C) and further Carboniferous and Permian clastics, 7 - effusive and extrusive products of basalt volcanism of Mesozoic evolution, locally with accompanying rocks.

